



NEW DATA ABOUT THE DISTRIBUTION AND ALTITUDINAL SPAN OF THE DALMATIAN RINGLET, *PROTEREBIA AFRA DALMATATA* (GODART, [1824]) (LEPIDOPTERA: SATYRINAE) IN CROATIA

Toni KOREN¹, Ivona BURIĆ², Ana ŠTIH³, Valerija ZAKŠEK⁴ & Rudi VEROVNIK⁴

¹Vela Traba 5, 52000 Pazin, Croatia; e-mail: koren.toni1@gmail.com

²Sinokoše 11, 21215 Kaštel Lukšić, Croatia

³Matije Divkovića 7, 10 000 Zagreb, Croatia

⁴University of Ljubljana, Biotechnical Faculty, Department of Biology, Večna pot 111, 1000 Ljubljana

Abstract - During the survey of butterfly fauna of Dalmatian mountains in the last years, *Proterebeia afra dalmata* (Godart, 1824) was found on several new sites, including the first records of this subspecies at higher altitudes (Poštak, Svilaja, Promina, Kamešnica and Dinara Mts.). In this paper we present a much more complete picture of the distribution of this presumably rare butterfly and discuss its altitudinal distribution. The species was found at altitude of almost 1500 m, therefore the characterization of the subspecies as a lowland butterfly can be dismissed.

KEY WORDS: *Proterebeia afra dalmata*, Dalmatia, distribution, altitudinal span

Izvleček – NOVI PODATKI O RAZŠIRJENOSTI IN RAZPONU NADMORSKIH VIŠIN POJAVLJANJA DALMATINSKEGA RJA VČKA, *PROTEREBIA AFRA DALMATATA* (GODART, [1824]) (LEPIDOPTERA: SATYRINAE) NA HRVAŠKEM

Med raziskavami favne metuljev dalmatinskih hribov v zadnjih letih smo vrsto *Proterebeia afra dalmata* (Godart, 1824) našli na več novih lokalitetah, vključno s prvimi najdbami te podvrste na višjih nadmorskih višinah (gore Poštak, Svilaja, Promina, Kamešnica in Dinara). V članku predstavljamo veliko popolnejšo sliko razširjenosti tega domnevno redkega metulja in razpravljamo o njegovi višinski razširjenosti. Vrsto smo našli na skoraj 1500 m višine, zato moramo zavreči opredelitev vrste kot nižinske.

KLJUČNE BESEDE: *Proterebeia afra dalmata*, Dalmacija, razširjenost, višinski razpon

Introduction

The genus *Proterebia* is monotypic with the distribution in Europe restricted to Greece (subspecies *Proterebia afra pyramus* de Louker & Dils, 1987), Russia south of the Urals (nominata subspecies), Crimea in Ukraine (subspecies *Proterebia afra krymaea* (Sheluzhko, 1929)) and Croatia (subspecies *Proterebia afra dalmata* (Godart, 1824)) (Tolman & Lewington 2008, Tshikolovets 2003).

P. afra dalmata is one of a few butterfly subspecies endemic to Croatia aside from *Erebia stirijs kleki* Lorković, 1955, *E. stirijs gorana* Lorković, 1985 and *E. gorge vagana* Lorković, 1955. Until 20 years ago its distribution in Croatia was very poorly known with records only from the surroundings of Zadar, Šibenik, Knin (Godart 1824; Stauder 1919–1927; Hafner 1994), and unknown locality on Korčula Island (Jakšić 1993). Over the past 10 years a lot of new data about the distribution of this species was published (Čelik *et al.* 2005; Mihoci & Šašić 2005, Zakšek, 2005; Mihoci & Šašić 2007) and eliminated many distribution gaps and expanded its known range from the Pag Island in the northwest to Biokovo Mts. in the South.

Dalmatian Ringlet flies from April to May in one generation (Tolman & Lewington 2008). *P. afra dalmata* is present predominantly in the Sub-Mediterranean region (with the exception of the unconfirmed record from the island of Korčula) (MIHOCI & ŠAŠIĆ, 2005). According to Horvatić (1971) the whole zone is characterized by unique climazonal vegetation of the association of *Quercus-Carpinetum orientalis croaticum*. Continental habitats of this species include overgrown submediterranean grassland community *Festuco-Koelerietum splendidis* or its progressive early succession stage with *Juniperus oxycedrus* on abandoned pastures (Čelik *et al.*, 2005).

While the Greek endemic subspecies *P. afra pyramus* is known to fly in the mountains ranging from an altitude of 600 m a.s.l. up to 1250 m a.s.l. (De Louker & Dils 1987; Abadijev 2002, Tolman & Lewington 2008) the Croatian subspecies, *P. afra dalmata* was believed to be confined to lower altitudes between 150 and 550 m a.s.l. (Tolman & Lewington 2008). According to Mihoci & Šašić (2005, 2007) the species flies up to 700 m a.s.l. (Donji Stublac, Biokovo Nature Park) and was found also almost at the sea level on island Pag (Zakšek 2005). Nevertheless the true altitudinal range of this subspecies was never a target to any systematic survey.

During the surveys of butterfly fauna of Dalmatia from 2005 onwards and targeted surveys of the Dalmatian mountains (Poštak, Prezid, Svilaja, Kamešnica, Dinara, Promina) in April and May 2010 *P. afra dalmata* was found on several new localities, including the first records of the subspecies at higher altitudes. In this paper we present new records of *P. afra dalmata* in Croatia and discuss conservation status, habitat preferences and altitudinal distribution of this subspecies.

Materials and methods

From the year 2005 onwards *P. afra dalmata* was found on 78 more or less distinct localities in central Dalmatia (Tab. 1, Fig. 1). Majority of these finds are from

new localities, only a few are confirmations of the known sites like the record from Vrelo Zrmanje, where it was found by Mihoci & Šašić (2007).

Specimens were caught by an entomological net, determined using Tolman & Lewington (2008) field guide, photographed and released on the same spot. Systematic follows Fauna Europaea (<http://www.faunaeur.org/>). For all known localities from the literature, an approximate altitude was determined using Google Earth tools, while for the new localities the altitude was determined using Garmin e-Trex vista GPS device.

Results

During surveys in the last years in central Dalmatia, *P. afra dalmata* was found on more than 60 new sites including the first records from five separate mountains: Poštak, Svilaja, Promina, Kamešnica and Dinara.

Additional records come from the Island of Pač, where the species seems to be widespread in open stony pastures from Novalja in the north to Povljan in the south. Despite similar suitable habitats on the mainland just south of the island and on nearby Vir Island, the species was not found there (Verovnik R., pers. observ.). Further southeast the butterfly is again common around Obrovac and further inland along Zrmanja and Krupa River valleys with isolated occurrence in Karišnica valley to the west. In this region the species was often observed in more humid habitats, on meadows along rivers, however the densities were mostly low, possibly indicating that these habitats are utilized by the species for occasional nectaring.

One of the main centers of the distribution of *P. afra dalmata* is the region SE of Zadar, around Benkovac, stretching as far south as hinterland of Šibenik. In this region the species was present almost in all suitable habitats, sometimes even entering the gardens in villages. Additional records confirm its contiguous presence also further south, especially around Primorski Dolac and as far as the slopes above Split and Trogir. The butterfly was particularly abundant on the ridge at Miljačka pass on 28.4.2007. Its distribution is probably more contiguous also further inland towards Driš, another center of the distribution of the species.

However, the most prominent new findings are from the mountain ranges in Dalmatia which are presented in more detail:

1. Svilaja is a mountain in Dalmatinska Zagora, parallel with the higher northern mountain range Dinara - Troglav. The highest peak of Svilaja is Bat (1508 m a.s.l.) (Poljak 2007). The large part of Svilaja, between Vrlika and Siverić is still covered with land-mines, so only some lower parts of the mountain are accessible and safe. On Svilaja, only a few specimens of *P. afra dalmata* were observed close to the main road at altitudes between 1050 and 1075 m. Butterflies were flying in typical habitat, on dry, grassy, sparsely bushy slopes partially covered with *Juniperus* bushes (Čelik et al. 2005) and were observed feeding on *Thymus* sp. and *Globularia* sp. plants.

2. Poštak is a mountain located south-east of Gračac. Its highest peak is Kučina Kosa (1441 m a.s.l.) (Poljak 2007). On Mt. Poštak *P. afra dalmata* was found near the highest peak at 1425 m a.s.l. and at the southern slopes at 1320 m a.s.l. Only few

specimens of *P. afra dalmata* were observed flying, but that could be attributed to the cloudy weather.

3. The Mt. Vela Promina is located west of Drniš. Unfortunately most of it is still covered with land-mines, so the only safe places to visit are the road to the peak and a small fenced area at the peak of Vela Promina, Čavinica. On Vela Promina *P. afra dalmata* was found near the road to the peak, on rocky slope covered with *Juniperus* bushes and on the peak itself, at 1148m. Only few specimens were observed on both locations.

4. The Dinara Mountain, with its peak Sinjal (1831 m a.s.l.), is the highest mountain in Croatia (Poljak 2007). It stretches from the northwest to southeast, between the Cetina River and Livanjsko polje. The southeastern part of the Dinara mountain range touches the Kamešnica Mountain. On Dinara *P. afra dalmata* was recorded in a continuous row of locations from the mountain hut Glavaš (561 m a.s.l.) to Martinova Košara (1300 m a.s.l.). It is interesting to note that *P. afra dalmata* was very abundant here and could be considered as a dominant species in suitable habitats. Due to bad weather conditions higher altitudes on the Dinara Mountain were not visited.

5. Mt. Kamešnica is located on the border with Bosnia and Herzegovina, north-east of Sinj. *P. afra dalmata* was recorded at three locations on the S side of the mountain at altitudes ranging from 780 to 1350 m a.s.l. in typical stony habitat.

Discussion

The new observations of *P. afra dalmata* fill many distribution gaps and possibly show the main shape of the area of its distribution in Croatia. The only prominent outlier, the record for Korčula Island without exact locality (Jakšić 1993), still remains to be confirmed. However, there are hardly any suitable habitats present on that island therefore the survey of the open stony areas along the main ridge of Pelješac Peninsula could prove much more rewarding. One of the unconfirmed localities remains Zadar, however this record could have been just a generalization of a find in a wide surrounding of the town (even as distanced as Benkovac region) or a genuine record from a site where the suitable habitats are now overgrown or destroyed by infrastructure. All other literature records are more or less in the vicinity of recent finds and well within the known distribution range (Fig. 1).

Recent finds only marginally extend the known range of the species further south-east to Župa and northwest on the Pag Island. Within the range the species is much more common and widespread as it was previously believed. So far the species has been recorded from more than 60 discrete sites excluding old records without information on exact locality. The main centers of distribution are around the Zrmanja River, region between Benkovac and Šibenik, and hills north of Split. The most prominent new records are from the mountains in the hinterland of Dalmatia where the species was found at higher altitudes than previously recorded.

The altitudinal span (Fig. 2) of the species is now much expanded with records above 1400 m. Occurrence of the species at such altitudes has for now been known only from Turkey (Hesselbarth et al. 1995) and its range in Central Asia (Tuzov et al. 1996), but also in Greece the records at altitudes of 1250 m (De Louker & Dils 1987)



Fig. 1. Distribution of Dalmatian Ringlet (*P. afra dalmata*) in Croatia. Grey spots represent old records, black spots represent new ones.




Fig. 2. Number of records of Dalmatian Ringlet (*P. afra dalmata*) in each of the altitudinal (divided into 100 m intervals for convenience).

visible during the surveys with low species abundances at sites with more advanced bush coverage.

New records clearly indicate how little we really know about the ecology of *P. afra dalmata* and how much we still have to learn. The large number of new records is also a good indication of the lack of systematic surveys due to limited number of researches and lack of funding. After “Distribution maps of the butterflies of Yugoslavia” (Jakšić 1988) no real effort was done to map the butterflies of Croatia, and produce a correct and modern distribution atlas of Croatian butterflies, so the distribution of most species in Croatia is still poorly known. Therefore we hope to

are of similar range. Currently the upper bound of the species in Croatia is at the peak of Mt. Poštak (1426 m a.s.l.); however its upper limit on Mt. Dinara was not explored and, given the suitable habitat, could be much higher. The majority of the records however are from lower altitudes, especially between 0 and 400 m. This can be easily explained by the fact that all but a few main distribution centers of the species are near the coast. This does not indicate the colline nature of *P. afra dalmata*, as the suitability of the habitat is possibly the only limiting factor of the distribution of the species. The distribution of the potentially suitable habitat which extends along the Adriatic coast and enters the Dalmatian mountains (Horvatić, 1971) is however much wider and additional surveys of suitable habitats elsewhere are much needed.

The habitats of the species observed during the field work varied between almost completely barren pastures on the Island of Pag, to almost entirely overgrown grasslands in the hinterland between Benkovac and Šibenik. Additionally, the species was found on montane grasslands and utilizing wet, flower rich meadows for nectaring. Despite this variability of habitats we fear that continuation of abandonment, resulting in conversion of grasslands into woods, could cause large scale local extinctions and thus contraction of the subspecies range. Such processes were clearly

trigger further research on the Dalmatian Ringlet and more generally, a concerted effort of mapping of butterfly distribution in Croatia.

Table 1. New records of the Dalmatian Ringlet (*Proterebia afra dalmata*) in Croatia. Localities are sorted in descending order from the highest a.s.l. to the lowest.

| Locality | Coordinates | | Alt. (m) | Date | Observer(s) |
|--|---------------|---------------|----------|------------------|--------------------|
| Poštak, Kučina Kosa | 44°15'18.67"N | 16°06'42.93"E | 1441 | May 19th, 2010 | Koren, Burić, Štiš |
| Kamešnica, Me ugorje | 43°41'58.44"N | 16°52'11.15"E | 1350 | May 24th, 2010 | Koren, Burić, Štiš |
| Dinara, Martinova kosara | 44°02'20.72"N | 16°24'39.04"E | 1300 | May 25th, 2010 | Koren, Burić, Štiš |
| Dinara, above Glavaš | 44°02'12.97"N | 16°24'53.41"E | 1175 | May 26th, 2010 | Koren, Burić, Štiš |
| Velika Promina, Cavinica | 43°55'27.49"N | 16°10'12.13"E | 1148 | May 23rd, 2010 | Koren, Burić, Štiš |
| Dinara, above Glavaš | 44°02'14.79"N | 16°25'11.03"E | 1130 | May 26th, 2010 | Koren, Burić, Štiš |
| Kamešnica, near the white road | 43°42'26.36"N | 16°51'15.01"E | 1085 | May 24th, 2010 | Koren, Burić, Štiš |
| Svilaja, above the main road | 43°51'27.89"N | 16°23'59.95"E | 1075 | May 20th, 2010 | Koren, Burić, Štiš |
| Dinara, above Glavaš | 44°01'22.83"N | 16°24'55.97"E | 925 | May 26th, 2010 | Koren, Burić, Štiš |
| Dinara, above Glavaš | 44°01'13.45"N | 16°01'13.45"E | 800 | May 26th, 2010 | Koren, Burić, Štiš |
| Kamešnica, Korita | 43°42'26.42"N | 16°48'24.68"E | 780 | May 24th, 2010 | Koren, Burić, Štiš |
| S slopes N from village Župa, Zagvozd | 43°20'15.10"N | 17°07'01.30"E | 758 | April 3rd, 2007 | Zakšek, V. |
| Vrelo Zрманje, near Marčetići | 44°13'58.74"N | 16°04'10.60"E | 722 | May 1st, 2010 | Koren, Burić, Štiš |
| Above village Otrić | 44°15'43.51"N | 16°02'42.51"E | 720 | May 25th, 2010 | Koren, Burić, Štiš |
| Dinara, Glavaš | 44°00'33.88"N | 16°25'01.65"E | 561 | May 26th, 2010 | Koren, Burić, Štiš |
| S of Drežnica, Mirlović polje, Dmiš | 43°48'45.80"N | 16°21'48.00"E | 552 | May 2nd, 2007 | Zakšek, V. |
| N from village Svaguše, Zagvozd, Grabovac | 43°25'06.70"N | 17°01'21.70"E | 485 | April 30th, 2007 | Zakšek, V. |
| E from village Milići, Lovrinčevići, Zagvozd | 43°22'57.90"N | 17°04'21.00"E | 473 | April 30th, 2007 | Zakšek, V. |
| Split, Gornje Kelami, on the pass north of the village east of Mt. Bituš | 43°37'54.08"N | 16°20'21.04"E | 459 | April 28th, 2007 | Verovnik, R. |
| W from G. Buljani, D. Bitelić, Hrvace, Sinj | 43°47'46.80"N | 16°35'50.70"E | 431 | April 29th, 2007 | Zakšek, V. |
| W of Janjići, Mirlović polje, Dmiš | 43°48'26.30"N | 16°20'40.80"E | 406 | May 2nd, 2007 | Zakšek, V. |
| NW from spring M. Rumin, Rumin, Hrvace, Sinj | 4°46'56.90"N | 16°39'06.20"E | 393 | April 29th, 2007 | Zakšek, V. |
| W of hill Lisnica, Sirovice, Dmiš | 43°50'47.30"N | 16°09'06.30"E | 383 | May 2nd, 2007 | Zakšek, V. |
| SW of G. Planjane, Unešić, Dmiš | 43°44'44.80"N | 16°11'43.80"E | 359 | May 2nd, 2007 | Zakšek, V. |
| SE of village Parati, D. Planjane, Dmiš | 43°46'54.30"N | 16°10'03.30"E | 340 | May 2nd, 2007 | Zakšek, V. |
| Ravni Golubići, near the pond | 44°11'47.40"N | 15°49'25.03"E | 340 | May 6th, 2010 | Koren, Burić, Štiš |
| Beneš Zрманja village | 44°09'45.71"N | 16°03'25.49"E | 320 | May 1st, 2010 | Koren, Burić, Štiš |
| Trogir, Prgomet, carstic plateau Muše on a small road from E side | 43°36'12.55"N | 16°13'13.78"E | 312 | April 28th, 2007 | Verovnik, R. |
| Vrelo Zрманje, Bogunovići | 44°12'12.31"N | 16°04'19.08"E | 303 | May 4th, 2010 | Koren, Burić, Štiš |
| Golubići, close to the Mt. Prezid | 44°06'19.68"N | 16°12'51.58"E | 298 | April 30th, 2010 | Koren, Burić, Štiš |
| W of hill Brušnjak, Višići, Sitno Donje, Primorski Dolac | 43°41'14.50"N | 16°09'44.80"E | 288 | May 2nd, 2007 | Zakšek, V. |
| Mokro polje | 44°03'29.07"N | 16°02'34.51"E | 286 | May 1st, 2010 | Koren, Burić, Štiš |
| W from Dirljić, Radošić, Trogir | 43°36'40.90"N | 16°16'27.80"E | 280 | May 2nd, 2007 | Zakšek, V. |
| Kaštel, Kozjak, on a small side road westwards to spring Češmenovac | 43°34'37.10"N | 16°19'26.55"E | 257 | April 28th, 2007 | Verovnik, R. |
| E of hill Vrljica, Bakovići, Primorski Dolac | 43°39'11.20"N | 16°12'07.90"E | 235 | May 2nd, 2007 | Zakšek, V. |
| Split, Radošić, along the road N of the crossroads | 43°36'27.83"N | 16°19'44.35"E | 234 | April 28th, 2007 | Verovnik, R. |
| S of hill Mrčelin umac, Sitno Donje, Primorski Dolac | 43°41'48.20"N | 16°09'22.70"E | 227 | May 2nd, 2007 | Zakšek, V. |
| Šibenik, Danilo, western part of the hill Baljčuš | 43°43'39.21"N | 15°59'06.83"E | 223 | April 28th, 2007 | Verovnik, R. |
| SE of Goriš, Konjevrate, Dmiš | 43°47'37.50"N | 16°01'54.50"E | 220 | May 2nd, 2007 | Zakšek, V. |
| Kaštel, Rudine, small valley with a road to Opor north of railroad | 43°34'18.73"N | 16°18'19.97"E | 202 | April 28th, 2007 | Verovnik, R. |
| W from hill Vjenac, Nadin, Benkovac | 44°07'55.56"N | 15°51'79.72"E | 202 | April 30th, 2005 | Zakšek, V. |
| Primorski Dolac, Troločke, on the plateau west of the village | 43°37'55.90"N | 16°14'30.12"E | 192 | April 28th, 2007 | Verovnik, R. |
| SW from village Mikulići, Benkovac | 44°01'19.80"N | 15°35'42.20"E | 176 | April 30th, 2005 | Zakšek, V. |
| Hills Kozarica and Kaštelina, Kašić, Pirovac | 43°52'07.90"N | 15°40'02.60"E | 161 | May 1st, 2005 | Zakšek, V. |
| N from village Raštevici, Polača, Benkovac | 44°03'01.00"N | 15°31'58.90"E | 156 | April 30th, 2005 | Zakšek, V. |
| SW of Gulin, Bilice, Šibenik | 43°46'31.70"N | 15°57'40.70"E | 154 | May 2nd, 2007 | Zakšek, V. |
| Obrovac, in the village Bogatnik at Šukare | 44°09'43.85"N | 15°49'13.01"E | 153 | March 3rd, 2008 | Verovnik, R. |
| W from hill V. Umac, Crljenik, Pirovac | 43°52'48.70"N | 15°40'54.60"E | 151 | May 1st, 2005 | Zakšek, V. |
| Krupa spring, close to the road | 44°11'46.67"N | 15°54'35.14"E | 143 | April 30th, 2010 | Koren, Burić, Štiš |

| Locality | Coordinates | | Alt. (m) | Date | Observer(s) |
|---|---------------|---------------|----------|------------------|--------------------|
| NW from village Begovići, Benovac | 44°02'26.80"N | 15°32'48.00"E | 142 | April 30th, 2005 | Zakšek, V. |
| Panići, near the road to vrelo Krupe | 44°10'46.45"N | 15°49'18.04"E | 133 | May 2nd, 2010 | Koren, Burić, Štih |
| Obrovac, along the road S of the town on the plateau | 44°11'41.90"N | 15°40'26.09"E | 132 | March 3rd, 2008 | Verovnik, R. |
| W of village Gornja Jagodnja, Polača, Benkovac | 44°00'52.00"N | 15°32'00.32"E | 129 | April 30th, 2005 | Zakšek, V. |
| SW from Komorovac, Novalja, Island of Pag | 44°34'00.00"N | 14°55'55.10"E | 127 | April 28th, 2005 | Zakšek, V. |
| W of Donje Miranje, Vrana, Pakoštane | 43°58'50.70"N | 15°34'59.70"E | 126 | April 30th, 2005 | Zakšek, V. |
| Hills Stubica and Drašnjak, Čubrići, Pirovac | 43°51'23.40"N | 15°41'28.30"E | 124 | May 1st, 2005 | Zakšek, V. |
| Sekulići, behind the cemetery | 44°13'04.57"N | 15°44'40.41"E | 120 | May 3rd, 2010 | Koren, Burić, Štih |
| W from village Zapužane, Benkovac | 44°00'55.50"N | 15°33'43.00"E | 118 | April 30th, 2005 | Zakšek, V. |
| E from village Ivkovići, Nadin, Škabrnja, Benkovac | 44°04'46.30"N | 15°46'30.15"E | 116 | April 30th, 2005 | Zakšek, V. |
| Obrovac, Krupa gorge, 2 km SE of the Krupa Monastery | 44°11'25.62"N | 15°52'28.34"E | 115 | March 3rd, 2008 | Verovnik, R. |
| S from village Čubrići, Pirovac | 43°51'30.00"N | 15°42'00.40"E | 114 | May 1st, 2005 | Zakšek, V. |
| Obrovac, along the road 0.5 km N of the turn for village Ribnica | 44°10'48.01"N | 15°37'02.61"E | 111 | March 3rd, 2008 | Verovnik, R. |
| Obrovac, Krupa, grasslands SE of the Krupa Monastery | 44°11'32.36"N | 15°53'09.82"E | 103 | March 3rd, 2008 | Verovnik, R. |
| E of village Ražnjevići, Polača, Benkovac | 44°01'10.10"N | 15°31'21.80"E | 101 | April 30th, 2005 | Zakšek, V. |
| Manastir, close to river Krupa | 44°11'26.33"N | 15°53'00.17"E | 101 | May 2nd, 2010 | Koren, Burić, Štih |
| Pastures between Kolansko blato and Kolan, Novalja, Island of Pag | 44°29'57.70"N | 14°56'20.50"E | 84 | April 27th, 2005 | Zakšek, V. |
| E from village Vidalići, Novalja, Island of Pag | 44°32'12.60"N | 14°57'47.80"E | 77 | May 2nd, 2005 | Zakšek, V. |
| Kaštel Žegarski, village | 44°09'24.14"N | 15°51'12.23"E | 75 | April 30th, 2010 | Koren, Burić, Štih |
| N of Smokvica, Poveljana, Island of Pag | 44°22'12.50"N | 15°12'00.20"E | 71 | April 30th, 2005 | Zakšek, V. |
| S from hill Bakrač, near Vrana lake, Pirovac | 43°51'58.20"N | 15°38'55.50"E | 67 | May 1st, 2005 | Zakšek, V. |
| N from marine Šimuni, Pag, Island of Pag | 44°28'28.60"N | 14°57'17.30"E | 65 | April 28th, 2005 | Zakšek, V. |
| N of hill Kosovac, Vrana, Pakoštane | 43°56'15.00"N | 15°34'20.70"E | 40 | April 30th, 2005 | Zakšek, V. |
| NW of Bošana, Pag, Island of Pag | 44°28'49.90"N | 15°00'03.80"E | 22 | April 28th, 2005 | Zakšek, V. |
| Turnić, S from Gorica, Poveljana, Island of Pag | 44°22'47.74"N | 15°07'16.52"E | 20 | April 28th, 2005 | Zakšek, V. |
| Stan, NW of Vlašići, Poveljana, Island of Pag | 44°21'25.70"N | 15°07'02.10"E | 16 | April 28th, 2005 | Zakšek, V. |
| Karin, Karišnica valley, middle part | 44°07'13.15"N | 15°37'43.30"E | 14 | March 3rd, 2008 | Verovnik, R. |
| Binjac, SE from Košljun, Pag, Island of Pag | 44°23'15.69"N | 15°06'14.04"E | 6 | April 28th, 2005 | Zakšek, V. |
| Drakovci, E from Poveljana, Island of Pag | 44°21'34.80"N | 15°08'15.40"E | 5 | April 28th, 2005 | Zakšek, V. |

References

- Abadjiev, S. P., 2002:** Types of Balkan butterflies in the collection of the Natural History Museum, London (Lepidoptera, Hesperioidea & Papilionoidea). *Neue Entomologische Nachrichten*, **53**: 3–53.
- Čelik, T., Zakšek, V., Vreš, B. & Verovnik, R., 2006:** Distribution and habitat characteristics of *Proterebia afra* (Fabricius, 1787) (Lepidoptera, Satyrinae) in Croatia. 1st Slovenian Entomological Symposium, Book of Abstracts, 8–9 pp.
- De Louker, S. & Dils, J., 1987:** The occurrence of *Proterebia phegea* Borkhausen in Greece with description of a new subspecies (Lepidoptera: Nymphalidae: Satyrinae). *Phegea* **15** (3): 157–160.
- Fauna Europaea (<http://www.faunaeur.org/>); accessed in October 2010.
- Hafner, I. 1994:** Verzeichnis der bei Knin gesammelten Schmetterlinge (Lepidoptera). *Natura Croatica*, **3** (2): 119–184.
- Hesselbarth, G., Van Oorschot, H. & Wagener, S., 1995:** Der Tagfalter der Türkei. Vol. 3., Selbstverlag S. Wagener, 847 pp., Bocholt.
- Horvatić, S., 1971:** Osnovne vegetacijske jedinice primorskog krša i pitanje njihove pojačane zaštite. Simpozij o zaštiti prirode u našem kršu. Odjel za prirodne nauke JAZU, 109–145. Zagreb.

- Jakšić, P.** 1988: Privremene karte rasprostranjenosti dnevnih leptira Jugoslavije (Lepidoptera, Rhopalocera). Jugoslavensko entomološko društvo. Posebna izdanja, 70 pp. Zagreb.
- Jakšić, P.** 1993: The M. Rogulja collection of the Rhopalocera (Lepidoptera) from the former state of Yugoslavia. *Entomologist's Gazette*, 44: 85–95.
- Godart, J. B.** 1824: Article Papillon. In: Latreille, P. A. & Godart, J. B. *Historie Naturelle. Entomologie Encyclopédie Méthodique*, 9: 530.
- Lorković, Z.**, 1955: Die Populations analyse zweier neuen stenochoren *Erebia* – Rassen aus Kroatien. *Biološki glasnik* 8: 53–76.
- Lorković, Z.**, 1985: Taxonomische Differenzierung der südöstlichsten Populationen von *Erebia stirijs* Godart 1824 (Lep., Satyridae). *Acta entomologica Jugoslavica* 21 (1–2): 9–15.
- Mihoci, I. & Šašić, M.**, 2005: New findings of the butterfly Dalmatian Ringlet, *Proterebia afra dalmata* (Godart, [1824]) (Lepidoptera, Satyrinae) in Croatia. *Natura Croatica* 14 (2): 121–129.
- Mihoci, I. & Šašić, M.**, 2007: New distribution data on the endemic butterfly *Proterebia afra dalmata* (Godart, [1824]) (Nymphalidae, Satyrinae) in Croatia. *Natura Croatica* 16 (3): 205–210
- Poljak, Ž.**, 2007: *Croatian mountains*, 4th edition (in Croat), Golden marketing-Tehnička knjiga, 142 pp. Zagreb
- Stauder, H.** 1919–1927: Die Schmetterlingsfauna der illyro-adriatischen Festland- und Inselzone (Faunula Illyrico Adriatica). *Ztschr. wiss. Insektenbiologie*.
- Tolman, T. & Lewington, R.** 2008: *Butterflies of Britain & Europe*. Harper Collins Publishers, 384 pp., London
- Tshikolovets, V. V.** 2003: *Butterflies of Eastern Europe, Urals and Caucasus*. Kyiv-Brno, 176 pp.
- Tuzov, V. K., Bogdanov, P. V., Devyatkin, A. L., Kaabak, L. V., Korolev, V. A., Murzin, V. S., Samodurov, G. D. & Tarasov, E. A.** 1997: *Guide to the butterflies of Russia and adjacent territories Volume 1*. Pensoft, Sofia-Moscow, 480pp.
- Zakšek, V.**, 2005: On the presence of *Proterebia afra* (Lepidoptera: Nymphalidae: Satyrinae) on the island of Pag, Croatia. *Phegea* 33 (3): 118–120.

Received / Prejeto: 16. 9. 2010